

**Amendments to the Specification:**

Please replace the paragraph beginning on page 21, line 21, with the following rewritten paragraph:

Preferred Fc variants include the following. In SEQ ID NO:2 (~~Figure 4~~Figures 3A and 3B) the leucine at position 15 may be substituted with glutamate; the glutamate at position 99, with alanine; and the lysines at positions 101 and 103, with alanines. In addition, one or more tyrosine residues can be replaced by phenylalanine residues.

Please replace the paragraph beginning on page 22, line 27, with the following rewritten paragraph:

A useful strategy for the PEGylation of synthetic peptides consists of combining, through forming a conjugate linkage in solution, a peptide and a PEG moiety, each bearing a special functionality that is mutually reactive toward the other. The peptides can be easily prepared with conventional solid phase synthesis (~~see, for example, Figures 5 and 6 and the accompanying text herein~~). The peptides are "preactivated" with an appropriate functional group at a specific site. The precursors are purified and fully characterized prior to reacting with the PEG moiety. Ligation of the peptide with PEG usually takes place in aqueous phase and can be easily monitored by reverse phase analytical HPLC. The PEGylated peptides can be easily purified by preparative HPLC and characterized by analytical HPLC, amino acid analysis and laser desorption mass spectrometry.